



AFRICAN SWINE FEVER (ASF)

DO YOU KNOW THE SIGNS AND SYMPTOMS OF THIS DISEASE?

CONTACT THE AUTHORITIES IMMEDIATELY IF YOU OBSERVE
ANIMALS SHOWING THESE SYMPTOMS

Ngaahi Faka'ilonga: The key clinical signs include:

- 'OKU MAFOLA 'A E MAHAKI NI 'I HA FA'AHINGA PE 'O E PUAKA TAUHI PE VAO 'I HA FA'AHINGA TA'U PE
This disease can affect both domestic and wild pigs of any age
- KULA/TAKA'ULI E KILI
Red/blotchy skin lesions
- MOFI LAHI 'O A'U KI HE MAFANA KO E 40°C - 42°C 'I HE HILI E 'AHO 'E 4-5 'ENE PIHIA HE MAHAKI
High fever 40°C - 42°C that begins 4-5 days after infection
- MALAVA KE FAKATOKANGA'I 'OKU NAU FAKAPUPUNGA MO NGALI 'OKU NAU FAKAMOKOSIA
Pigs may huddle together and are usually shivering
- TA'ETA'E FIE KAI
Loss of appetite
- MANAVA TAMAKI PEA TALETALE
Abnormal breathing and coughing
- VAIVAIA PEA 'IKAI LAVA KE FA'A TU'U KI 'OLUNGA
Lethargic-refuses to stand
- LUA PEA/PE FAKALELE MO E TU'U TOTO
Vomiting and/or diarrhea with bloody discharge
- HAFU E MATA MO E IHU
Heavy discharge from eyes and nose
- TAMATO, FA'ELE TA'EHOKO, VAIVAIA E 'UHIKI 'OKU MALAVA KE HOKO
Miscarriage, abortions, still births and weak litters can occur



©INIA-CISA/Carmina Gallardo, except top. ©IZS-UM